Open Journal of Biomedical Research

Open J Biomed Res

REVIEW ARTICLE

Microbial Endocrinology: The Influence of the Gut Microbiome on Metabolic and Endocrine Health

Abdurraouf Zaet[1, 2, 3] *, Amel M. Aswehlee[4], Karema Zeat[2,5]

[1]Department of Dental Technology, Faculty of Medical Technology, University of Zawia, Libya.
[2]Biomedical Research Team, University of Zawia, Libya.
[3]Libyan Society for Biomedical Research, Libya.
[4]Digitodontics International Academy, France.
[5]Department of Medical Nutrition, Faculty of Medical Technology, University of Zawia, Libya.

ABSTRACT

Purpose: The human gastrointestinal tract harbors a complex microbial ecosystem that transcends its traditional digestive role, functioning as a dynamic endocrine organ capable of producing bioactive signaling molecules. This emerging paradigm, termed microbial endocrinology, recognizes that gut microorganisms both respond to and synthesize neuroendocrine hormones, establishing bidirectional communication pathways with host physiological systems.

Materials and Methods: This review systematically examines the interplay between intestinal microbiota and host endocrine networks, with particular emphasis on elucidating mechanistic pathways through which microbial-derived metabolites, neurotransmitters, and immune mediators govern metabolic homeostasis, stress physiology, thyroid function, and reproductive health.

Results: Gut microbes demonstrate substantial capacity to modulate host energy metabolism, with fecal microbiota transfer from obese donors inducing adiposity in recipient animals. Dysbiotic configurations correlate with metabolic syndrome, type 2 diabetes, polycystic ovary syndrome, and thyroid autoimmunity through mechanisms including short-chain fatty acid signaling, lipopolysaccharide-mediated inflammation, bile acid transformation, and enterohepatic hormone circulation. Therapeutic modulation via probiotics, prebiotics, and fecal transplantation shows promise but requires personalization due to substantial inter-individual heterogeneity in microbial composition.

Conclusion: The intestinal microbiome operates as an integral component of host endocrine architecture, with dysbiosis representing a modifiable risk factor for metabolic and hormonal pathologies. Precision microbiome interventions tailored to individual endocrine profiles represent a transformative frontier for managing metabolic syndrome, diabetes, and endocrine disorders. *(Open J Biomed Res 2024;3:15-20)*

Keywords

Gut microbiome, Microbial endocrinology, Metabolic syndrome, Type 2 diabetes, Short-chain fatty acids. Hypothalamic-pituitary-adrenal axis.

AUDIOABSTRACT

ARTICLE INFO

***Corresponding Author:**

Dr. Abdurraouf Zaet

Department of Dental Technology, Faculty of Medical Technology, University of Zawia, Libya.

Email: a.zaet@zu.edu.ly

Received: 5 May 2024
Accepted: 17 June 2024
Published: 20 June 2024

INTRODUCTION

The conceptual framework of microbial endocrinology, pioneered by Mark Lyte, fundamentally reconceptualizes host-microbe interactions by establishing that microorganisms actively participate in neuroendocrine signaling.[1] Rather than serving as passive inhabitants of the gastrointestinal tract, bacteria demonstrate sophisticated capacity to detect, produce, and respond to hormonal and neurochemical signals including catecholamines and other bioactive compounds challenging historical boundaries between microbial and host physiology.[1,2]

The scale of this microbial metabolic capacity warrants emphasis: the human gut accommodates approximately 10^14^ microbial cells encoding genetic machinery that outnumbers host genes by two orders of magnitude.[3] This extensive enzymatic repertoire enables microbiota to function as a genuine endocrine organ, synthesizing and modulating hormones, neurotransmitters, and signaling metabolites that enter systemic circulation.[4] Consequently, dysbiosis characterized by disrupted microbial diversity and composition has emerged as a central pathogenic mechanism underlying metabolic syndrome, type 2 diabetes mellitus, obesity, polycystic ovary syndrome, and thyroid dysfunction.

Articles published in the Open J Biomed Res are licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

THE GUT MICROBIOME AS AN ENDOCRINE ORGAN

Microbial Production of Hormone-like Molecules

Intestinal bacteria generate diverse bioactive compounds that directly interface with host endocrine physiology. Short-chain fatty acids primarily acetate, propionate, and butyrate arise from bacterial fermentation of dietary fibers and function as critical signaling ligands. These metabolites engage G protein-coupled receptors (GPR41, GPR43) on enteroendocrine L-cells, triggering secretion of glucagon-like peptide-1 (GLP-1) and peptide YY (PYY) hormones that govern insulin release, satiety signaling, and energy balance.[5]

Clinical investigations demonstrate that targeted colonic delivery of propionate enhances insulin sensitivity and reduces hepatic lipid accumulation, confirming direct metabolic benefits beyond mere energy substrate provision.[6]

Beyond short-chain fatty acids, commensal organisms actively metabolize steroid hormones. The aggregate genetic capability of microbiota to process estrogens termed the "estrobolome" modulates circulating hormone levels through β-glucuronidase-mediated deconjugation, thereby influencing estrogen receptor activation in distal tissues.[7] This microbial hormone processing represents a fundamental mechanism linking intestinal ecology to systemic endocrine function.

Microbial Neurotransmitter Synthesis

Intestinal bacteria synthesize neuroactive compounds capable of crossing the blood-brain barrier or activating vagal neural pathways. Specific *Lactobacillus* and *Bifidobacterium* strains produce γ-aminobutyric acid (GABA), the principal inhibitory neurotransmitter, while *Escherichia*, *Bacillus*, and *Saccharomyces* generate norepinephrine, serotonin, and dopamine.[8]

Remarkably, the gastrointestinal tract produces approximately 95% of bodily serotonin, with microbiota regulating this synthesis through tryptophan metabolism modulation and enterochromaffin cell stimulation.[9] This microbial contribution to neurochemical milieu establishes the foundation for microbiota-gut-brain axis communication with profound implications for both metabolic regulation and psychological wellbeing.

GUT MICROBIOME AND METABOLIC HEALTH

Obesity and Energy Homeostasis

Gut microbiota substantially influences host energy extraction and adipose tissue accumulation. Gnotobiotic mice receiving microbiota from obese human donors exhibit greater weight gain and fat deposition compared to those colonized with lean donor material demonstrating causal involvement in obesity pathogenesis.[10] This enhanced energy harvest stems from microbial degradation of otherwise indigestible polysaccharides coupled with modulation of host lipid metabolism gene expression.[11]

Individuals with obesity typically display reduced microbial diversity and altered *Bacteroidetes-Firmicutes* proportional representation, though precise taxonomic signatures remain under active investigation.[12] Microbial metabolites directly impact metabolic pathways: lipopolysaccharide from Gram-negative organisms triggers low-grade systemic inflammation a hallmark of obesity and insulin resistance through Toll-like receptor 4 activation,[13] whereas butyrate enhances insulin sensitivity and skeletal muscle mitochondrial function.[14]

Type 2 Diabetes Mellitus

Dysbiotic configurations strongly associate with type 2 diabetes mellitus, marked by depletion of butyrate-producing taxa such as Faecalibacterium prausnitzii alongside expansion of opportunistic pathogens.[15] Metagenomic profiling reveals distinct functional alterations in diabetic microbiomes, including diminished short-chain fatty acid biosynthetic capacity and enhanced oxidative stress resistance machinery.[16]

Microbiota additionally govern bile acid metabolism, converting primary to secondary bile acids that signal through farnesoid X receptor and Takeda G protein-coupled receptor 5 mechanisms that regulate glucose homeostasis and insulin sensitivity.[17] The therapeutic efficacy of metformin partially derives from microbiome restoration toward eubiosis, suggesting that intestinal microbial modulation may constitute a primary pharmacological mechanism for this widely prescribed antidiabetic agent.[18]

MICROBIAL ENDOCRINOLOGY OF MAJOR HORMONAL AXES

The Hypothalamic-Pituitary-Adrenal (HPA) Axis

Intestinal microbiota profoundly modulates stress reactivity and glucocorticoid regulation. Gnotobiotic animals display exaggerated HPA axis responses to restraint stress, with elevated corticosterone and adrenocorticotropic hormone effects reversible through specific bacterial

colonization.[19] Chronic psychological stress reciprocally alters microbiome composition, increasing Clostridium abundance while reducing overall diversity, thereby compromising intestinal barrier integrity and promoting systemic inflammation.[20]

Microbial metabolites modulate HPA axis tone through multiple pathways: butyrate strengthens intestinal barrier function and reduces lipopolysaccharide translocation and subsequent inflammatory HPA activation.[21] Probiotic interventions using Lactobacillus rhamnosus demonstrate anxiolytic effects and cortisol normalization in both preclinical models and human trials.[22,23]

Thyroid Axis

Accumulating evidence implicates intestinal microbiota in thyroid hormone regulation. Dysbiosis correlates with reduced triiodothyronine (T3) concentrations in experimental models, potentially through microbial modulation of deiodinase enzymatic activity. The gut-thyroid axis operates through several convergent mechanisms: bacterial β-glucuronidase influences hormone enterohepatic circulation; lipopolysaccharide-induced inflammation impairs thyroid hormone receptor expression; and short-chain fatty acids modulate immune tolerance relevant to autoimmune thyroiditis.[24]

Hypothyroid patients exhibit characteristic microbiome signatures with reduced Bifidobacterium and Lactobacillus representation, while autoimmune thyroid disease associates with increased intestinal permeability and dysbiotic configurations.[25] Probiotic supplementation demonstrates preliminary efficacy in improving thyroid function parameters and reducing antibody titers, warranting expanded clinical investigation.[26]

Reproductive Hormones

The estrobolome critically influences estrogen-dependent pathologies. Microbial β-glucuronidase deconjugates estrogenic compounds in the intestine, enhancing their enterohepatic recirculation and systemic bioavailability.[27] Dysbiosis featuring elevated β-glucuronidase activity associates with estrogen-driven conditions including breast malignancy, endometriosis, and polycystic ovary syndrome. Women with polycystic ovary syndrome display characteristic microbial alterations including reduced diversity and diminished short-chain fatty acid-producing bacteria correlating with hyperandrogenism and insulin resistance.[28]

Testosterone metabolism similarly undergoes microbial modification. Intestinal bacteria metabolize testosterone and its precursors, with antibiotic-induced dysbiosis altering circulating testosterone concentrations and impacting reproductive function in animal models.[29] This bidirectional relationship between sex steroids and intestinal microbiota represents a crucial dimension of microbial endocrinology with implications spanning reproductive health and metabolic disease.

THE MICROBIOTA-GUT-BRAIN AXIS IN ENDOCRINE REGULATION

The vagus nerve serves as the principal neural conduit transmitting intestinal signals to central neuroendocrine circuits. Vagal afferents express receptors for short-chain fatty acids, hormones, and cytokines, conveying microbial information to hypothalamic and brainstem nuclei.[30] This communication influences appetite, metabolism, mood, and stress responses integrating endocrine and neurological functions into unified physiological networks.

Microbial regulation of tryptophan metabolism provides another critical interface. Intestinal organisms' direct tryptophan toward either kynurenine or serotonin synthetic pathways, profoundly affecting peripheral and central serotonin availability.[31] This metabolic diversion influences psychological states while simultaneously modulating glucose homeostasis through serotonin-mediated insulin secretion and hepatic glucose production regulation. The convergence of microbial, neural, and endocrine signaling underscores the holistic nature of physiological coordination.

THERAPEUTIC IMPLICATIONS AND CLINICAL APPLICATIONS

Probiotics and Prebiotics

Strategic microbiome manipulation offers substantial therapeutic promise. Akkermansia muciniphila supplementation improves metabolic parameters in overweight and obese individuals reducing insulin resistance and inflammatory markers.[32] Prebiotic fibers that selectively nourish beneficial taxa similarly enhance glycemic control and satiety hormone secretion.[5] Current challenges include identifying strain-specific effects and optimizing delivery systems to ensure durable colonization and functional metabolic activity.

Fecal Microbiota Transplantation

Fecal microbiota transplantation represents the most direct microbiome intervention. In metabolic syndrome, transplantation from lean donors transiently enhances insulin sensitivity, though response durability varies considerably.[33] Success

depends upon donor microbiota engraftment and recipient baseline composition, emphasizing personalized therapeutic approaches.[34] Ongoing investigations explore applications for polycystic ovary syndrome and additional endocrine disorders, though long-term safety and efficacy require further validation.

Dietary Interventions

Dietary patterns fundamentally sculpt microbiome composition and endocrine output. Mediterranean dietary patterns rich in polyphenols and fiber promote short-chain fatty acid-producing bacteria while reducing inflammatory potential, correlating with improved metabolic and endocrine biomarkers.[35] Conversely, high-fat diets rapidly induce dysbiosis and metabolic endotoxemia within days, demonstrating remarkable microbiome responsiveness to nutritional change.[36] Time-restricted feeding and intermittent fasting beneficially remodel microbiome composition, enhancing circadian metabolic alignment.[37]

CHALLENGES AND FUTURE DIRECTIONS

Despite considerable progress, substantial obstacles persist. Establishing causality versus association remains challenging in human investigations, while tremendous inter-individual microbiome variability complicates universal therapeutic recommendations.[38] The field requires standardized analytical protocols, enhanced functional gene expression characterization beyond taxonomic profiling, and predictive modeling for treatment stratification.

Future research must address temporal dynamics of microbiome-endocrine interactions particularly during developmental critical periods and aging transitions. Multi-omics integration with detailed phenotyping will enable precision microbiome medicine tailored to individual metabolic and endocrine profiles. Understanding cumulative environmental impacts including pollutants, antibiotics, psychological stress on microbiome endocrine function represents an essential priority for public health strategies.

CONCLUSION

Microbial endocrinology provides an integrative framework for comprehending how intestinal bacteria influence host metabolic and endocrine health. The microbiome functions as a pervasive endocrine organ, generating hormones, neurotransmitters, and metabolites that integrate with host physiology through diverse signaling pathways. Dysbiosis contributes to metabolic syndrome, diabetes, and endocrine disorders pathogenesis, while targeted microbiome interventions offer novel therapeutic avenues. Recognizing the microbiome as a central node within endocrine networks will prove essential for developing next-generation treatments for metabolic and endocrine diseases.

REFERENCES

1. Lyte M. Microbial endocrinology in the microbiome-gut-brain axis: how bacterial production and utilization of neurochemicals influence behavior. PLoS Pathog. 2013;9:e1003726. doi: 10.1371/journal.ppat.1003726.
2. Lyte M. Microbial endocrinology: Host-microbiota neuroendocrine interactions influencing brain and behavior. Gut Microbes. 2014;5:381-9. doi: 10.4161/gmic.28682.
3. Clarke G, Stilling RM, Kennedy PJ, Stanton C, Cryan JF, Dinan TG. Minireview: Gut microbiota: the neglected endocrine organ. Mol Endocrinol. 2014;28:1221-38. doi: 10.1210/me.2014-1108.
4. Tremaroli V, Bäckhed F. Functional interactions between the gut microbiota and host metabolism. Nature. 2012;489:242-9. doi: 10.1038/nature11552.
5. Canfora EE, Jocken JW, Blaak EE. Short-chain fatty acids in control of body weight and insulin sensitivity. Nat Rev Endocrinol. 2015;11:577-91. doi: 10.1038/nrendo.2015.128.
6. Chambers ES, Viardot A, Psichas A, Morrison DJ, Murphy KG, Zac-Varghese SE, et al. Effects of targeted delivery of propionate to the human colon on appetite regulation, body weight maintenance and adiposity in overweight adults. Gut. 2015;64:1744-54. doi: 10.1136/gutjnl-2014-307913.
7. Plottel CS, Blaser MJ. Microbiome and malignancy. Cell Host Microbe. 2011;10:324-35. doi: 10.1016/j.chom.2011.10.003.
8. Strandwitz P. Neurotransmitter modulation by the gut microbiota. Brain Res. 2018;1693(Pt B):128-133. doi: 10.1016/j.brainres.2018.03.015.
9. Yano JM, Yu K, Donaldson GP, Shastri GG, Ann P, Ma L, et al. Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis. Cell. 2015;161:264-76. doi: 10.1016/j.cell.2015.02.047.
10. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. Nature. 2006;444:1027-31. doi: 10.1038/nature05414.
11. Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, Nagy A, et al. The gut microbiota as an environmental factor that regulates fat storage. Proc Natl Acad Sci U S A. 2004;101:15718-23. doi: 10.1073/pnas.0407076101.

12. Finucane MM, Sharpton TJ, Laurent TJ, Pollard KS. A taxonomic signature of obesity in the microbiome? Getting to the guts of the matter. PLoS One. 2014;9:e84689. doi: 10.1371/journal.pone.0084689.
13. Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, et al. Metabolic endotoxemia initiates obesity and insulin resistance. Diabetes. 2007;56:1761-72. doi: 10.2337/db06-1491.
14. Gao Z, Yin J, Zhang J, Ward RE, Martin RJ, Lefevre M, et al. Butyrate improves insulin sensitivity and increases energy expenditure in mice. Diabetes. 2009;58:1509-17. doi: 10.2337/db08-1637.
15. Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. Nature. 2012;490, 55–60. https://doi.org/10.1038/nature11450.
16. Karlsson FH, Tremaroli V, Nookaew I, Bergström G, Behre CJ, Fagerberg B, et al. Gut metagenome in European women with normal, impaired and diabetic glucose control. Nature. 2013;498:99-103. doi: 10.1038/nature12198.
17. Wahlström A, Sayin SI, Marschall HU, Bäckhed F. Intestinal Crosstalk between Bile Acids and Microbiota and Its Impact on Host Metabolism. Cell Metab. 2016;24:41-50. doi: 10.1016/j.cmet.2016.05.005.
18. Wu H, Esteve E, Tremaroli V, Khan MT, Caesar R, Mannerås-Holm L, et al. Metformin alters the gut microbiome of individuals with treatment-naive type 2 diabetes, contributing to the therapeutic effects of the drug. Nat Med. 2017;23:850-858. doi: 10.1038/nm.4345.
19. Sudo N, Chida Y, Aiba Y, Sonoda J, Oyama N, Yu XN, et al. Postnatal microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in mice. J Physiol. 2004;558:263-75. doi: 10.1113/jphysiol.2004.063388.
20. Bailey MT, Dowd SE, Galley JD, Hufnagle AR, Allen RG, Lyte M. Exposure to a social stressor alters the structure of the intestinal microbiota: implications for stressor-induced immunomodulation. Brain Behav Immun. 2011;25:397-407. doi: 10.1016/j.bbi.2010.10.023.
21. Kelly JR, Kennedy PJ, Cryan JF, Dinan TG, Clarke G, Hyland NP. Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related psychiatric disorders. Front Cell Neurosci. 2015;9:392. doi: 10.3389/fncel.2015.00392.
22. Bravo JA, Forsythe P, Chew MV, Escaravage E, Savignac HM, Dinan TG, et al. Ingestion of Lactobacillus strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. Proc Natl Acad Sci U S A. 2011;108:16050-5. doi: 10.1073/pnas.1102999108.
23. Messaoudi M, Lalonde R, Violle N, Javelot H, Desor D, Nejdi A, et al. Assessment of psychotropic-like properties of a probiotic formulation (Lactobacillus helveticus R0052 and Bifidobacterium longum R0175) in rats and human subjects. Br J Nutr. 2011;105:755-64. doi: 10.1017/S0007114510004319.
24. Fröhlich E, Wahl R. Microbiota and Thyroid Interaction in Health and Disease. Trends Endocrinol Metab. 2019;30:479-490. doi: 10.1016/j.tem.2019.05.008.
25. Knezevic J, Starchl C, Tmava Berisha A, Amrein K. Thyroid-Gut-Axis: How Does the Microbiota Influence Thyroid Function? Nutrients. 2020;12:1769. doi: 10.3390/nu12061769.
26. Talebi S, Karimifar M, Heidari Z, Mohammadi H, Askari G. The effects of synbiotic supplementation on thyroid function and inflammation in hypothyroid patients: A randomized, double blind, placebo controlled trial. Complement Ther Med. 2020;48:102234. doi: 10.1016/j.ctim.2019.102234.
27. Kwa M, Plottel CS, Blaser MJ, Adams S. The Intestinal Microbiome and Estrogen Receptor-Positive Female Breast Cancer. J Natl Cancer Inst. 2016;108:djw029. doi: 10.1093/jnci/djw029.
28. Torres PJ, Siakowska M, Banaszewska B, Pawelczyk L, Duleba AJ, Kelley ST, et al. Gut Microbial Diversity in Women With Polycystic Ovary Syndrome Correlates With Hyperandrogenism. J Clin Endocrinol Metab. 2018;103:1502-1511. doi: 10.1210/jc.2017-02153.
29. Jakobsson HE, Jernberg C, Andersson AF, Sjölund-Karlsson M, Jansson JK, Engstrand L. Short-term antibiotic treatment has differing long-term impacts on the human throat and gut microbiome. PLoS One. 2010;5:e9836. doi: 10.1371/journal.pone.0009836.
30. Bonaz B, Bazin T, Pellissier S. The Vagus Nerve at the Interface of the Microbiota-Gut-Brain Axis. Front Neurosci. 2018;12:49. doi: 10.3389/fnins.2018.00049.
31. Agus A, Planchais J, Sokol H. Gut Microbiota Regulation of Tryptophan Metabolism in Health and Disease. Cell Host Microbe. 2018;23:716-724. doi: 10.1016/j.chom.2018.05.003.
32. Depommier C, Everard A, Druart C, Plovier H, Van Hul M, Vieira-Silva S, et al. Supplementation with Akkermansia muciniphila in overweight and obese human volunteers: a proof-of-concept exploratory study. Nat Med. 2019;25:1096-1103. doi: 10.1038/s41591-019-0495-2.
33. Vrieze A, Van Nood E, Holleman F, Salojärvi J, Kootte RS, Bartelsman JF, et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. Gastroenterology. 2012;143:913-6.e7. doi: 10.1053/j.gastro.2012.06.031.
34. Li SS, Zhu A, Benes V, Costea PI, Hercog R, Hildebrand F, et al. Durable coexistence of donor and recipient strains after fecal microbiota transplantation. Science. 2016;352:586-9. doi: 10.1126/science.aad8852.

35. Singh RK, Chang HW, Yan D, Lee KM, Ucmak D, Wong K, et al, Liao W. Influence of diet on the gut microbiome and implications for human health. J Transl Med. 2017;15:73. doi: 10.1186/s12967-017-1175-y.
36. Martinez-Guryn K, Hubert N, Frazier K, Urlass S, Musch MW, Ojeda P, et al. Small Intestine Microbiota Regulate Host Digestive and Absorptive Adaptive Responses to Dietary Lipids. Cell Host Microbe. 2018;23:458-469.e5. doi: 10.1016/j.chom.2018.03.011.
37. Zarrinpar A, Chaix A, Yooseph S, Panda S. Diet and feeding pattern affect the diurnal dynamics of the gut microbiome. Cell Metab. 2014;20:1006-17. doi: 10.1016/j.cmet.2014.11.008.
38. Franzosa EA, Morgan XC, Segata N, Waldron L, Reyes J, Earl AM, et al, Huttenhower C. Relating the metatranscriptome and metagenome of the human gut. Proc Natl Acad Sci U S A. 2014;111:E2329-38. doi: 10.1073/pnas.1319284111.